

AD-A060 675


NAVAL AEROSPACE MEDICAL RESEARCH LAB DETACHMENT MICHG--ETC F/G 6/3  
BIOELECTRIC INDICANTS OF DIVER'S ABILITY TO PERFORM USEFUL WORK--ETC(U)  
MAR 78 R S KENNEDY

UNCLASSIFIED

TM-78-01


NL

1 OF 1  
ADA  
060675



END  
DATE  
FILMED

1 -79  
DDC



NATIONAL BUREAU OF STANDARDS  
MICROCOPY RESOLUTION TEST CHART

**LEVEL***12 nu*


6 Bioelectric Indicators of Diver's Ability to Perform Useful Work.

9 Technical memo.

Presented at the  
Undersea Medical Society Workshop  
Bethesda, Maryland

11 March 1978

12 20p.



10 Robert S. Kennedy PhD  
~~CDR USN USN~~

Naval Aerospace Medical Research Laboratory Detachment  
New Orleans, Louisiana

This document has been approved  
for public release and sale; its  
distribution is unlimited.

NAVAL AEROSPACE MEDICAL RESEARCH LABORATORY DETACHMENT  
NEW ORLEANS, LOUISIANA

The opinions are those of the author and do not necessarily reflect those  
of the Department of the Navy.

391

221

78 10 20 050

*Gu*

AD A060675

DDC FILE COPY

# ABSTRACT

Presently, the majority of working dives are performed in shallow depths and the performances which are expected of the diver emphasize physical rather than mental work. Nevertheless, dives are continuing to be made to deeper depths and in the future, increasing demands will be made of the diver for information processing and decision making. It was hypothesized that a neglected indicant of an operator's ability to perform useful work in air is eye movement activities. It was felt that aspects of eye movements could provide a useful index of the level of information processing or mental load of a diver and secondly, that research into eye movements might aid us in understanding better the effects of the physical environment experienced by a diver.

The present paper reports recent research in air between eye movements and performance. The findings are positive and offer promise that spectral analysis of eye movement velocities could provide valid early indication of behavioral dysfunction in compressed gas and under water.

REFERENCE TO	
OPS	WHI Section <input checked="" type="checkbox"/>
SEC	DOT Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
NOTIFICATION	
BY	
BY NAME AND TELEPHONE CODES	
DATE	
TIME	
A	

78 10 20 050



## INTRODUCTION

The present paper will discuss the relationship between certain characteristics of eye movement and overall central nervous system status, beginning with the study of vestibularly induced nystagmus.

The relationship of eye movement to vestibular stimulation is well known and has been described many times (Kennedy 1972). Specifically, as the head moves to a new position, the eyes lag behind the skull and effectively permit one to continue looking at the starting point. This is not a passive response due to inertia nor is it due to visual fixation, although both of those factors are present. Rather, the slow compensatory deviation of the eyes in the skull (which serves to keep the eyes fixed over the earth) - is caused directly by a vestibular signal and occurs also in the dark. This eye movement is the slow phase of nystagmus and has as its probable site of origin the vestibular nuclei (Gernandt 1959). The second phase - the fast phase - is considered to be compensatory to the slow.

In Guedry's (1965) psychophysiological review he referenced about 20 papers where the subject's mental state modified recorded vestibular nystagmus. Gernandt (1959) has suggested that the fast phase is influenced by the reticular activating system (RAS) and Wolfe (1966) has retrieved habituated nystagmus by RAS stimulation. Further, it was shown that the fast phase was absent in patients who lacked a pontine reticular formation, (Daroff & Hoyt 1971). Therefore, it was decided to expose a large group of people to vestibular stimulation while recording nystagmus and measuring their performance on a vigilance task. The hypothesis had been that vigilance performance would bear some relationship to the quality of the fast phase. In other words, knowledge of the fast phase could be predictive of the vigilance performance.

Vigilance test - three tones which were clearly audible and distinguishable were presented randomly for an hour. The subject's task was to monitor the low (8 pulses/minute) and middle (6 pulses/minute) and ignore the high (5 pulses/minute). Each occurrence of a tone was mentally counted and a key was pressed when a given tone had been sounded four times. The subject then began again for that tone. The subject's score was percent correct for each 5/minute period. Figure 1 shows the temporal distribution of tones and Figure 2 shows the expected values for six versions of this test. The one labelled "two channel complex" is described above and was employed in the eye movement experiment.

Nystagmus - the subject was dark adapted for 20 minutes to partially control for fluctuations in the cornea-retinal potential (Kennedy 1972), and was then exposed to sinusoidal oscillation about the spinal axis for 50 minutes. Stimulus parameters were

a 75° displacement every five seconds. (.2 Hz; 46.8/sec peak velocity and 53.50/sec<sup>2</sup> peak angular acceleration.)

Lateral eye movements were obtained by standard electro-oculographic techniques with electrodes at the outer canthi. Pre and post calibrations were not significantly different. One hundred fifty healthy student pilots comprised the experimental population, 50 were tested for vigilance only, 50 for nystagmus only and 50 for both.

Scoring - the procedure used is best described as an examination of the fast phase. In general, 100% of a cycle with good nystagmus received a 10, and almost no nystagmus was scored 1.0. Two lower categories were used, .5 and .1 for finer determinations. The reliability of the method was good. ( $r = .95$ ). It should be noted that while using mainly fast phase (presence/absence) in scoring, Wendt (1965) felt that other eye movements qualify as habituation, particularly with shorter arcs of oscillation.

## RESULTS

The next slide (Figure 3) shows the performance of these three groups. The vigilance performance of both groups was not significantly different but the quality of nystagmus in the group without the vigilance task decayed more rapidly than for those who did mental work. This is what one would expect from the literature as reported by Guedry (1965). Vigilance and nystagmus follow the same time course ( $r = .93$ ).

In order to determine whether the vigilance performance of a particular subject could be predicted from his nystagmus at any point in time, each subject's scores were correlated within a session. The average of these correlations was  $r = .30$  and significant ( $P < .0001$ ). Another correlation was obtained for the 500 matched scores (50 subjects and 10 five minute time frames). This operation is statistically indefensible but has a practical utility for vestibular investigators. For instance, you may wish to predict a person's vigilance score regardless of who he is or what time in the session it is. This correlation is  $r = .49$  ( $P < .001$ ).

In conclusion, this researcher feels that under the experimental conditions used, a general correspondence exists between the quality of the fast phase and the alertness as measured by a vigilance task. This correspondence supports what would be expected from the neurophysiological literature regarding the pathways of the fast phase and has both applied and scientific implications. It suggested that (1) perhaps the fast phase of vestibular nystagmus could be used as a quantitative and independent index of arousal,



so that a person's performance in a job could be monitored without interfering with his work; (2) perhaps other quick flicks of the eyes (optokinetic fast phase, involuntary movements and microsaccades) reflect similar mechanisms. If true, they also could provide indicants of the level of arousal. If the latter could be shown to be so, it might be possible for a computer to analyze the eye movements of a pilot or diver and when they meet a criterion level of drowsiness, inform him of this condition and remove him from the situation. Because of these results, the second phase of this experiment was begun, wherein no vestibular stimulation at all was used, but merely the subjects sitting in the dark, performing the same two channel monitoring task and again recording movements but with some variation. Two recording techniques were employed: first, surface electrode as before with EEG type electrodes at the outer canthi and secondly on infrared scleral reflection device. The comparisons of these two methods will be reported elsewhere and need not be discussed further, except to indicate that each has different characteristics which would make it more or less feasible, depending on the working conditions, diving conditions, etc. Secondly, an analysis of eye movement data which differed from before was undertaken. It was felt that the reason for the previously obtained relationship between eye movement and performance occurred because something happened relative to the fast phase - whereas the slower eye movements remained essentially the same; albeit with a slight change in phase and in gain.

In a previous study the data were all scored by hand and an inspection of the data revealed that the presence of a brisk, strong, fast phase contributed to alert scores and what appeared like slower fast phases occurred when vigilance was poor. It was hypothesized that the briskness or velocity of the fast phase of eye movements may be generally indicative of alertness or of overall potential for work. Therefore, a new method for scoring eye movements was employed: specifically, a spectral analysis of eye movement velocities which were sampled 100x/sec, directly from the infra-red eye movement recording device was performed. These were totalized by computer and printed out as scores each minute. The next Figure (4) shows the data for the entire experiment. Note that nearly all the eye movement power is found in the range 0 - 100°/sec. Figure 5 shows the same type of information but in this case early and late eye movement are separated for the various ranges of velocities. It may be seen that the 100° - 200°/sec category is changed so that there are fewer "later" than "earlier" and the converse occurs with 0 - 100°/sec eye movements. It should be emphasized that while these differences are small, they are also highly reliable, therefore likely to be real. It should also be noted that these differences, and other data to be presented, are consistent within subjects as well. The next figure (6) shows the eye movement data for a single channel performance - the simplest test comparing early and late. It may be seen that later eye

movements show less activity in the  $100^{\circ} - 200^{\circ}/\text{sec}$  category. Additionally, vigilance performance also degraded on this test. Figure 7 shows two things; (1) the same effect, but note that the average scores (approximately .89 versus .75) correspond to overall differences in the task loading (see also Table I). Figure 8 completes the picture where a similar effect is present viz., more overall eye movement between  $100^{\circ} - 200^{\circ}/\text{sec}$ , with a greater task load, and also a drop in performance with time on task. All these data are compiled in the next slide, Fig. 9. What has occurred, is that different task loads produce different distributions of eye movement velocities whereby higher loads appear to coincide with more rapid movements. Secondly, as performance degrades with time on task, there appears to be a corresponding drop in eye movement activity but this relationship is less clear cut because the least amount of reduction occurred in the two channel test; concurrently there was also very little drop in performance. The overall speculation is twofold: when larger numbers of fast eye movements are available, the operator may have greater potential for information load; secondly, operator workload is characterized by the lower number of eye movements in the  $0 - 100^{\circ}/\text{sec}$  range, whereby relatively more of them indicate a light load. In general, these two factors would be correlated most of the time, but not necessarily. The eye movement data of the present study would tend to support this notion, particularly when considered in connection with the percent correct scores on the various forms of the counting tests. In summary, in all cases with time on task there is a change toward relatively more "slow" ( $0 - 100^{\circ}/\text{sec}$ ) later and relatively more "fast" ( $100^{\circ} - 200^{\circ}/\text{sec}$ ) early. Secondly, the eye movement level early in a test is consonant with this finding, whereby the overall amount of fast activity (see Table I) is proportional to the level of workload.

In addition to the above it is felt that the technology necessary for eye movement measurement might also bring with it assessment monitoring techniques that would have the following additional spinoff for the working diver:

First, untoward vestibular effects which result in nystagmus can be recorded.

Second, information about where a diver is looking can be recorded for purposes of display, design and other training of diver work operations.

Third, eye movements may be considered to be a cortical evoked response indicator, and changes thereto could be monitored.

Fourth, the recording technology would promote development of visually coupled systems like those used in aircraft, viz., a) the visual target acquisition system (VTAS) for remote handling of mass and b) the energy management display which enables the pilot to better utilize his resources.



In conclusion, the analysis of velocity of eye movements can be a useful, sensitive and reliable parameter which may be predictive of changes in the state of an individual. This research could progress from a hyperbaric chamber to open sea conditions in two to three years.



TABLE I

FREQUENCY OF EYE MOVEMENT VELOCITIES DURING AUDITORY MONITORING  
FOR 6 SUBJECTS (BEATS/SEC)

	<u>ALL TESTS</u>	<u>EARLY IN SESSION</u>	<u>LATE IN SESSION</u>
	0-100 DPS	5.12	5.25
	100-200 "	.85	.78
	200-300 "	.20	.19
I CHANNEL	0-100 DPS	5.25	5.45
	100-200 "	.75	.65
	200-300 "	.19	.17
II CHANNEL	0-100 DPS	4.90	5.00
	100-200 "	.89	.85
	200-300 "	.26	.27
III CHANNEL	0-100 DPS	5.20	5.40
	100-200 "	.94	.80
	200-300 "	.15	.14

## REFERENCES

- Gernandt, B., Vestibular mechanisms, In: Field and Magoun (Eds), Handbook of Physiology. Phila: Williams & Wilkins, Vol. 1, 1959, 549-564.
- Guedry, F.E., Psychophysiological studies of vestibular function. In: Neff (Ed) Contributions to sensory physiology v.e. 1965, Academic Press: N.Y. pp 63-135.
- Daroff, R.B. & Hoyt, W.F., In: Bach-y-Rita & Collins (Eds), Symposium on the Control of Eye Movements 1971, Academic Press, N.Y.
- Kennedy, R.S., The relationship between habituation to vestibular stimulation and vigilance: Individual differences and subsidiary problems. PhD dissertation, University of Rochester 1972. NAMRL Monograph No. 20, July 1972. (Also published in Dissertation Abstracts International, No. 72-78, 764, November 1972).
- Wendt, G.R., The nature of adaptation to oscillatory rotation. In: The role of the vestibular organs in the exploration of space, 1965, NASA special publication number SP-77, pp 133-139.
- Wolfe, J.W., Evidence for cerebellar control over habituation of vestibular nystagmus, Unpublished PhD dissertation, University of Rochester, 1966.

## LIST OF FIGURES

- FIG. 1      Five minute swatch of temporal distribution of tones
- FIG. 2      Performance on six different versions of a vigilance task
- FIG. 3      Nystagmus and vigilance in three different groups
- FIG. 4      Spectral density of eye movement velocities over all six subjects
- FIG. 5      A comparison of early and late eye movement velocity categories in all vigilance tasks combined
- FIG. 6      A comparison of early and late eye movement velocity categories in the one channel vigilance task
- FIG. 7      A comparison of early and late eye movement velocity categories in the two channel vigilance task
- FIG. 8      A comparison of early and late eye movement velocity categories in the three channel vigilance task
- FIG. 9      A comparison of early and late eye movement velocity categories on three different vigilance tasks
- FIG. 10     Performance in percent correct on three different vigilance tasks



TONES



# AUDITORY VIGILANCE TASK





FIGURE 2. A COMPARISON OF PERFORMANCES ON SIX AUDITORY VIGILANCE TASKS



FIGURE 3. VESTIBULAR NYSTAGMUS AND VIGILANCE FOR THREE GROUPS OF NAVAL AVIATION PERSONNEL



FREQUENCY OF VELOCITY




FIGURE 4. SPECTRAL DENSITY OF EYE MOVEMENT VELOCITIES  
OVER SIX SUBJECTS






FIGURE 5. A COMPARISON OF EARLY AND LATE EYE MOVEMENT  
VELOCITY CATEGORIES IN ALL VIGILANCE TASKS COMBINED




FIGURE 6. A COMPARISON OF EARLY AND LATE EYE MOVEMENT  
VELOCITY CATEGORIES IN THE ONE CHANNEL VIGILANCE  
TASK






FIGURE 7. A COMPARISON OF EARLY AND LATE EYE MOVEMENT  
VELOCITY CATEGORIES IN THE TWO CHANNEL VIGILANCE  
TASK






FIGURE 8. A COMPARISON OF EARLY AND LATE EYE MOVEMENT  
VELOCITY CATEGORIES IN THE THREE CHANNEL VIGILANCE  
TASK




FIGURE 9. A COMPARISON OF EARLY AND LATE EYE MOVEMENT VELOCITY CATEGORIES ON THREE DIFFERENT VIGILANCE TASKS



COMBINED PERFORMANCE  
OF ALL SUBJECTS




FIGURE 10. PERFORMANCE IN PERCENT CORRECT ON  
THREE DIFFERENT VIGILANCE TASKS